

## Recombinant Bovine FGF2

**Product Part No.:** FGF2-221P

**Product quantity:** 50 µg, 100 µg, 500 µg, 1 mg

### Meet Your Product

<b>Product Details</b>	FGF2 is a member of the FGF family (one of 23). <sup>1</sup> Bovine FGF2 is a single, non-glycosylated, polypeptide chain containing 155 amino acids with molecular mass of 17.3 kDa. The FGF family of proteins drive broad mitogenic and cell survival activities including cell proliferation, differentiation, survival, and apoptosis. <sup>2</sup> FGF2 binds to the FGF receptor (FGFR) members, and activates downstream signaling proteins, like ERK1/2. <sup>345</sup>
<b>Synonyms</b>	Fibroblast growth factor basic; Heparin-binding growth factor 2 (HBGF2); Prostatropin
<b>Sequence (monomer)</b>	>sp P03969 FGF2_BOVINE MAAGSITTLPALPEDGGSGAFPPGHFKDPKRLYCKNGGFFLRIHPDGRVDGVREKSDP HIKLQLQAEERGVSISIKGVCANRYLAMKEDGRLLASKCVTDECFFFERLESNNYNTYR SRKYSSWYVALKRTGQYKLGPKTGPGQKAILFLPMSAKS
<b>Formulation</b>	Lyophilized protein from 0.22 µm filtered solution in PBS, pH 7.4
<b>Purity &amp; Identity</b>	Verified by SDS PAGE to be >95% pure
<b>Endotoxin</b>	Below threshold of <1 EU/µg
<b>Protein Content</b>	Concentration verified by total protein assay
<b>Biological Activity</b>	Determined by proliferation of NIH-3T3 cells and binding to receptor via SPR
<b>Application</b>	Research Use Only. Not for diagnostic or therapeutic applications

### Storage & Handling

Products are lyophilized and shipped at ambient temperature. Please follow the storage and handling instructions below after receiving the product.

<b>Storage</b>	Store at 5 ± 3°C upon receipt.
<b>Stability</b>	Stable as supplied for 12 months from date of manufacturing.
<b>Reconstitution</b>	Gently tap down the vial to ensure that all lyophilisate is collected at the bottom of the vial. Reconstitute the product in PBS to at least 100 µg/mL by gently pipetting the solution down the sides of the vial. Avoid vigorous shaking that can cause foaming and protein denaturation. Keep on ice. Aliquot and store at 5 ±



	3°C for up to 2 weeks or at < -20°C for up to 3 months. Avoid repeated freeze-thaw cycles by aliquoting reconstituted products.
<b>Recommendation</b>	As an additional precaution, after adding reconstituted products to media, filter-sterilize before use in cell culture.

## Frequently Asked Questions

### 1. Are there any cross-species activity with this growth factor?

Yes, bovine FGF2 shares 99% sequence homology with human FGF2; expected cross reactivity against human, rat, porcine, mouse, chicken, horse and other mammalian systems.

### 2. Why can't I see a pellet in the vial?

Lyophilized powder may not be visible for many reasons including, but not limited to, dislodged powder being stuck on the cap or lack of carrier proteins that make the product difficult to see. Please tap or centrifuge the vial to bring all the material down to the bottom and reconstitute the product as outlined above.

### 3. Are there any stability concerns with freeze thawing?

Repeated freeze thawing is not recommended, as this may damage the protein products resulting in reduced functionality. After reconstitution, please aliquot into suitable sizes for one time use and freeze.

### 4. Can I add BSA as a carrier?

Yes. Adding a carrier protein like BSA (0.1%) enhances protein stability, increases shelf-life, and allows the recombinant protein to be stored at a more dilute concentration.

### 5. Is this protein bioactive?

Yes, this protein was tested for activity on its cognate receptor via SPR, as well as proliferation of NIH3T3 cells.

## References

1. Xie Y, Su N, Yang J, Tan Q, Huang S, Jin M, Ni Z, Zang B, Zang D, Luo F, Chen H, Sun X, Feng J, Chi L and Chen L. FGF/FGFR signaling in health and disease. *Signal Transduct Target Ther.* 2020 Sep 2;5(1):181.
2. Bikfalvi A, Klein S, Pintucci G, Rifkin DB. Biological roles of fibroblast growth factor-2. *Endocr Rev.* 1997 Feb;18(1):26-45.
3. Dai S, Zhou Z, Chen Z, Xu G, Chen Y. Fibroblast growth factor receptors (FGFRs): structures and small molecule inhibitors. *Cells.* 2019 Jun 18;8(6).
4. Kimelman D, Kirschner M. Synergistic induction of mesoderm by FGF and TGF-beta and the identification of an mRNA coding for FGF in the early *Xenopus* embryo. *Cell.* 1987 Dec 4;51(5):869-77.
5. Kang HB, Kim JS, Kwon H-J, Nam KH, Youn HS, Sok D-E, Lee Y. Basic fibroblast growth factor activates ERK and induces c-fos in human embryonic stem cell line MizhES1. *Stem Cells Dev.* 2005 Aug;14(4):395-401.