



Product Data Sheet - Recombinant Human Prolactin, Tag Free

Product details	
Product Name	: Recombinant Human Prolactin, Tag Free
Product Part Number	: PRL-111
Product Description	<p>Human Prolactin, also known by its gene name PRL, is a hormone produced by the pituitary gland. It is primarily responsible for promoting lactation in the mammary glands, but it also has pleiotropic effects in both males and females¹⁻⁶. Prolactin typically exists in two forms: a glycosylated and a non-glycosylated version⁶. Each prolactin molecule is believed to bind with two PRL-receptor molecules⁷. In addition to its role in lactation, peripheral prolactin production is associated with breast and prostate cancer development, regulation of reproductive function, and immune system control. Recent research has suggested that PRL has cytoprotective effects on pancreatic islet cells and liver tissue, which could be crucial for the success of transplantation in clinical settings⁸⁻¹⁰.</p> <p>In physiology, PRL appears in both glycosylated and non-glycosylated forms in specific ratios that are dynamic based on other factors; furthermore, their affinity to receptor and half-life <i>in vivo</i> are determined by post translational modifications.</p>
Synonyms	Mammotropin, Luteotropic hormone, Luteotropin, PRL
Sequence	<div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;"> Human Prolactin (Met 1 - Cys 227) P01236 * </div> <p style="text-align: center;">*SELEVLFQ</p> <p>There is a short 8 amino acid overhang post-cleavage. Full sequence available upon request. Total Amino acids: 235 aa. Calculated MW: 26.8 kDa.</p>
Formulation	Lyophilized protein in 50 mM Tris-HCl, 100 mM NaCl, pH 7.4
Purity & Identity	≥90% pure, Verified by SDS PAGE. Identity has been verified during characterization. Data available upon request.
Safety	Endotoxin: <1 EU/μg. Mycoplasma: Negative
Protein Content	Concentration verified by Total protein assay.
Biological Activity	Determined by a cell proliferation assay using Nb2-11 cell line and binding to the receptor via SPR. Effective concentration 50 (EC ₅₀) < 3



	ng/ml.
Application	Research Use Only. Not for diagnostic or therapeutic applications.

Storage & Handling

Shipping	Products are lyophilized and shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperatures below.
Storage	Upon arrival, store the lyophilized protein at -20°C.
Stability	Stable as supplied for 3 months from retest date. For continued use beyond the retest date, contact the manufacturer.
Reconstitution	Gently tap down the vial to ensure that all lyophilisate is collected at the bottom of the vial. Reconstitute the product in sterile water to at least 100 µg/mL by gently pipetting the solution down the sides of the vial. Avoid vigorous shaking that can cause foaming and protein denaturation. Keep on ice. Following reconstitution, aliquot and store at < -70°C for up to 3 months under sterile conditions. Avoid repeated freeze-thaw cycles.
Recommendation	As an additional precaution, after adding reconstituted products to media, filter-sterilize before use in cell culture.

Frequently Asked Questions

- Are there any cross-species activity with this growth factor?**
 Yes, even though Human Prolactin shares approximately 60% to 65% similarity with rat and mouse prolactin, it can still trigger prolactin receptors from other species³.
- Why can't I see a pellet in the vial?**
 Lyophilized powder may not be visible for many reasons including, but not limited to, dislodged powder being stuck on the cap or lack of carrier proteins that make the product difficult to see. Please tap or centrifuge the vial to bring all the material down to the bottom and reconstitute the product as outlined above.
- Are there any stability concerns with freeze thawing?**
 Repeated freeze thawing is not recommended, as this may damage the protein products resulting in reduced functionality. After reconstitution, please aliquot into suitable sizes for one time use and freeze.



- **Can I add BSA as a carrier?**

Yes. Adding a carrier protein like BSA (0.1%) enhances protein stability, increases shelf-life, and allows the recombinant protein to be stored at a more dilute concentration

- **Is this protein bioactive?**

Yes, this protein was tested for activity on proliferation of NB2-11 cells.

References

1. Cooke NE, Coit D, Shine J, Baxter JD, Martial JA. Human prolactin. cDNA structural analysis and evolutionary comparisons. *J Biol Chem.* 1981 Apr 25;256(8):4007–16.
2. Grattan DR, Kokay IC. Prolactin: a pleiotropic neuroendocrine hormone. *J Neuroendocrinol.* 2008 Jun;20(6):752–63.
3. Ben-Jonathan N, LaPensee CR, LaPensee EW. What can we learn from rodents about prolactin in humans? *Endocr Rev.* 2008 Feb;29(1):1–41.
4. Forbes K, Westwood M. Maternal growth factor regulation of human placental development and fetal growth. *J Endocrinol.* 2010 Oct;207(1):1–16.
5. Goffin V, Hoang DT, Bogorad RL, Nevalainen MT. Prolactin regulation of the prostate gland: a female player in a male game. *Nat Rev Urol.* 2011 Oct 4;8(11):597–607.
6. Price AE. Studies on the microheterogeneity and in vitro activity of glycosylated and nonglycosylated recombinant human prolactin separated using a novel purification process. *Endocrinology.* 1995 Nov 1;136(11):4827–33.
7. Broutin I, Jomain J-B, Tallet E, van Agthoven J, Raynal B, Hoos S, et al. Crystal structure of an affinity-matured prolactin complexed to its dimerized receptor reveals the topology of hormone binding site 2. *J Biol Chem.* 2010 Mar 12;285(11):8422–33.
8. Wailemann RA, Terra LF, Oliveira TC, Dos Santos AF, Gomes VM, Labriola L. Heat shock protein B1 is required for the prolactin-induced cytoprotective effects on pancreatic islets. *Mol Cell Endocrinol.* 2018 Dec 5;477:39–47.
9. Yamamoto T, Ricordi C, Mita A, Miki A, Sakuma Y, Molano RD, et al. beta-Cell specific cytoprotection by prolactin on human islets. *Transplant Proc.* 2008 Mar;40(2):382–3.
10. Ostróžka-Cieślík A, Dolińska B. Pharmacological Benefits and Risk of Using Hormones in Organ Perfusion and Preservation Solutions in the Aspect of Minimizing Hepatic Ischemia-Reperfusion Injury during Storage. *Biomed Res Int.* 2019 Nov 11;2019:6467134.

Copyright © 2024 All rights reserved by Future Fields Cellular Agriculture and Research Ltd. FUTURE FIELDS® is a trademark of Future Fields Cellular Agriculture and Research Ltd., a company registered in Canada.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. Purchasers must determine the suitability of the product(s) for their particular use.

End of Product Data sheet